

# Swine Feed Efficiency: Influence of Temperature

**Author**  
Tom G. Miller, Iowa State University

**Reviewers**  
Mike D. Tokach, Kansas State University  
Joel M. DeRouchey, Kansas State University

## Introduction

Pig performance, including feed efficiency, is maximized when animals are kept within their thermoneutral zone. When describing the environmental conditions that the animal is raised in, it is important to realize that there are several factors influencing pig comfort. “Effective” environmental temperature is a better measure than “air” temperature, as it takes into account relative humidity, air movement (velocity), flooring type and condition, insulation of the building, evaporative cooling, and huddling of pigs. Pigs of different size and age have different thermoneutral comfort zones, as younger, smaller pigs have higher temperature requirements and have a smaller tolerance for temperature changes (Figure 1). As pigs grow, their thermoneutral zone expands and they are able to tolerate a much wider range of temperature before growth rate and feed efficiency are affected.

## Effects of Environmental Temperature on Performance

Pigs are homoeothermic animals maintaining body temperature within narrow limits under varying environmental conditions, balancing heat loss and heat production. When temperatures fall below the pig’s lower critical limit, the pig must produce additional heat; they may do so by increasing energy intake or diverting more energy from growth to maintenance, or a combination of the two. In finishing pigs fed a balanced diet, feed efficiency mostly depends on the level of energy intake, maintenance requirement, and the efficiency with which ME above maintenance is used for growth. Changing feed (energy) intake is one way the animal is able to cope with changes in temperature. During times of cold stress, a pig will increase feed intake. During times of heat stress, feed intake is reduced.




Figure 1. A model illustrating effective temperatures for pigs of different weights.

The effects of temperature outside of grow-finish pigs’ thermoneutral temperature zone are shown in Table 1. Pigs consume more feed and are less efficient at converting feed to gain when the effective temperature is below their lower critical limit as more nutrients are used for maintenance and less are available for growth. When temperatures are above the upper critical limit, pigs are less active and consume less feed; however, their maintenance requirements may still be relatively high due to increased respiration rate. Respiration rates increase dramatically during heat stress because pigs lack the ability to sweat and, therefore, panting is a means to remove body heat. Additionally, the animal’s average daily feed intake is lower which slows average daily gain resulting in more days to market and impacting feed efficiency.

Table 1. Effect of environmental temperature on the performance of growing pigs (50 to 125 pounds)<sup>1</sup>

Item	Temperature, °F		
	50	72.5	95
ADG, lb	1.70	1.76	1.41
ADFI, lb	4.86	4.21	3.52
Feed/gain	2.91	2.41	2.52

<sup>1</sup>Summary of 3 experiments; Stahly and Cromwell, 1979, 1981.

## Temperature and Feed Intake

Table 2 shows the increased feed intake of different sized pigs for each degree Fahrenheit below their lower critical temperature (LCT). Example: A finisher pig weighing 220 lb which is kept 10 °F cooler than its LCT would be expected to eat 0.43 lb more feed per day than a pig housed at the correct temperature. In a 2,400 head finishing barn, daily consumption of feed would be increased by over 1000 lb/day or 3.5 tons per week.

Table 2. Increase in feed intake when the effective environmental temperature falls below the pigs' lower critical temperature (LCT)

Category	Weight, lb	Increase in Feed Intake, lb/d
<b>Piglet</b>	4	0.005
<b>Nursery</b>	20	0.008
<b>Nursery</b>	45	0.016
<b>Grower</b>	100	0.026
<b>Finisher</b>	220	0.043
<b>Thin Sow</b>	313	0.073
<b>Heavier Sow</b>	>450	0.042

\*Based on a diet ME of 2900 kcal/kg or 1315 kcal/lb. Data adapted from Holmes and Close, 1977

## Additional Key Considerations

- Manipulating the amount of fiber in the diet can be used as a management tool whether in a hot or cold environment as heat generation is greater with increasing dietary fiber.
- Reducing crude protein or increasing diet energy through fat addition can improve performance during heat stress.
- Ventilation management is vital to providing the correct air conditions, preventing cold, damp or hot, humid environmental conditions.
- Evaporative cooling (drillers, misters, or foggers) is an effective tool to reduce thermal stress and the effective environmental temperature felt by the pig during hot weather.
- Genetic lines with lower residual feed intake may have an improved ability to cope with heat stress.

## References

- Baker, J.E. 2004. Effective environmental temperature. *J Swine Health Prod.* 12(3):140-143.
- Holmes C.W. and W.H. Close. 1977. The influence of climatic variables on energy metabolism and associated aspects of productivity in the pig. p 51-73 in *Nutrition and the climatic environment*. Butterworths, London, UK.
- McFarlane J.M. How do your pigs really feel? 2004. Marysville, Ohio: Animal Environment Specialists.
- Renaudeau, D., H. Glibert, and J. Noblet. 2012. Effect of climatic environment on feed efficiency in swine. p183 –210 in: *Feed Efficiency of Swine*. Wageningen, Netherlands.
- Stahly, T.S. and G.L. Cromwell. 1979. Effect of environmental temperature and dietary fat supplementation on the performance and carcass characteristics of growing and finishing swine. *J. Anim. Sci.* 49:1478-1488.
- Stahly, T.S., G.L. Cromwell, and J.R. Overfield. 1981. Interactive effects of season of year and dietary fat supplementation, lysine source and lysine level on performance of swine. *J. Anim. Sci.* 53:1269-1277.

... and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue SW, Washington, DC 20250-9410, or call 800-795-3272 (voice) or 202-720-6382 (TDD). USDA is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Cathann A. Kress, director, Cooperative Extension Service, Iowa State University of Science and Technology, Ames, Iowa